JOURNAL OF HEALTH, MEDICAL RESEARCH AND INNOVATIONS

VOLUME: 02 ISSUE: 03 (2024)

https://www.jhmri.com

Research Article

# MENTAL HEALTH CHALLENGES AMONG HEALTHCARE WORKERS DURING PANDEMIC OUTBREAKS: A SYSTEMATIC REVIEW

***Dr. Usman Khalid***

*Institute of Public Health, Dow University of Health Sciences, Karachi, Pakistan.*

## Abstract

*The COVID-19 pandemic, among other health crises, has significantly strained healthcare systems worldwide, especially impacting the mental well-being of frontline workers. This systematic review investigates the psychological challenges faced by healthcare workers (HCWs) during pandemic outbreaks, including anxiety, depression, burnout, post-traumatic stress, and insomnia. We reviewed studies published between 2000 and 2024, focusing on pandemics such as SARS, MERS, Ebola, and COVID-19. Data were extracted from databases like PubMed, Scopus, and Google Scholar. The results indicate a high prevalence of psychological distress among HCWs, exacerbated by factors such as long working hours, fear of infection, lack of protective equipment, and insufficient institutional support. The findings underscore the urgent need for structured psychological interventions and institutional resilience mechanisms to support healthcare providers in future outbreaks.*

**Keywords:** *Mental Health, Healthcare Workers, Pandemic Stress, Psychological Resilience.*

## Introduction

Pandemic outbreaks such as SARS (2002), MERS (2012), Ebola (2014–2016), and most notably COVID-19 (2019–2023), have not only challenged the global health infrastructure but have also exposed frontline healthcare workers to substantial psychological strain [1][2]. HCWs are often required to perform under extreme pressure, facing high infection risks, long duty hours, ethical dilemmas, and emotional fatigue [3][4]. The convergence of these stressors has led to widespread mental health disorders among medical personnel, with studies reporting increased rates of anxiety (up to 44%), depression (up to 37%), and insomnia (up to 36%) during COVID-19 [5][6].

Despite growing recognition of these challenges, there remains a lack of systematic synthesis that consolidates findings across different pandemic scenarios. This article aims to fill that gap by conducting a comprehensive review of existing literature to identify patterns, risk factors, and potential interventions to mitigate mental health impacts on HCWs.

## 1. Methodology and Data Sources

This systematic review followed the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines [7] to ensure a transparent and reproducible selection process. The primary aim was to synthesize available evidence on the mental health challenges faced by healthcare workers (HCWs) during various pandemic outbreaks.

### Search Strategy:

A comprehensive literature search was conducted across three major scientific databases: PubMed, Scopus, and Web of Science, covering publications from January 2000 to March 2024. The search terms included combinations of:

- "mental health" OR "psychological impact" OR "stress" OR "anxiety"
- AND "healthcare workers" OR "frontline workers"
- AND "pandemic" OR "COVID-19" OR "SARS" OR "MERS" OR "Ebola"

### Inclusion Criteria:

- Peer-reviewed empirical studies (quantitative, qualitative, or mixed-methods)
- Focused on mental health outcomes among HCWs during pandemics
- Included clearly defined psychological outcomes (e.g., anxiety, depression, PTSD, burnout)
- Studies available in English language
- Studies conducted in any geographical context, with subgroup analysis for South Asia and Pakistan

### Exclusion Criteria:

- Non-peer-reviewed content (e.g., opinion pieces, editorials, news articles)
- Articles without empirical psychological outcome measures
- Studies unrelated to pandemics (e.g., general stress among HCWs in routine settings)
- Duplicate records or studies with insufficient methodological clarity

### Screening and Selection Process:

Titles and abstracts were screened independently by two reviewers. Full texts of potentially eligible articles were retrieved for detailed review. Disagreements were resolved through discussion or a third reviewer.

### Data Extraction and Quality Appraisal:

A standardized data extraction form was developed to collect the following variables:

- Author(s), year, country
- Pandemic type
- Study design and sample characteristics
- Type and prevalence of mental health outcomes
- Risk/protective factors identified

The Critical Appraisal Skills Programme (CASP) checklist [8][9] was used to assess the quality and methodological rigor of included studies. Studies scoring below 6 out of 10 on the CASP scale were excluded to ensure credibility of evidence synthesis.

**Final Sample:**

Out of 1,542 initial records, 96 studies met the inclusion criteria after full-text screening and quality appraisal.

**2. Psychological Outcomes and Prevalence**

Pandemic outbreaks have significantly affected the psychological well-being of healthcare workers (HCWs), manifesting in various mental health conditions. Across the studies reviewed, the most frequently reported psychological outcomes include anxiety**,** depression**,** post-traumatic stress disorder (PTSD)**,** burnout**,** and sleep disturbances [10].

**Anxiety and Depression**

Anxiety was the most prevalent condition reported, particularly during the COVID-19 pandemic, with rates reaching up to 45% among nurses and 38% among physicians, attributed to fears of contagion, overwhelming workloads, and isolation from families [11]. Similarly, depression affected nearly 30% of frontline workers, with symptoms including low mood, fatigue, and hopelessness. Studies highlighted that younger HCWs and those with fewer years of experience were especially vulnerable to depressive symptoms [11].

**Burnout and Occupational Stress**

Burnout—characterized by emotional exhaustion, depersonalization, and reduced personal accomplishment—was notably high among ICU and emergency department staff, who faced frequent patient deaths, ethical dilemmas, and triage decisions [12]. A meta-analysis showed burnout prevalence ranging from 32% to 60%**,** depending on specialty and workload intensity.

**Post-Traumatic Stress and Insomnia**

PTSD symptoms were commonly reported during the early phases of pandemics, particularly among workers involved in high-risk areas such as infectious disease units and emergency care. Rates of PTSD symptoms during the SARS and COVID-19 outbreaks ranged from 10% to 25% [10]. Sleep disorders, including insomnia, affected up to 36% of HCWs, often exacerbated by shift work, psychological strain, and lack of recovery time.

**Gender and Role-Based Differences**

Research consistently revealed gender disparities, with female HCWs experiencing significantly higher levels of anxiety and depression than their male counterparts. This may be due to a combination of occupational exposure and domestic caregiving responsibilities [13]. Additionally, nurses reported higher stress levels than physicians, likely due to closer patient interactions and longer shifts without breaks.

**Prevalence Summary Table (From Meta-Analysis):**

| Disorder | Prevalence (%) | Population | Reference |
|---|---|---|---|
| **Anxiety** | 30–45% | Nurses & Physicians | [11] |
| **Depression** | 20–30% | Frontline HCWs | [11] |
| **Burnout** | 32–60% | ICU, ER Staff | [12] |
| **PTSD Symptoms** | 10–25% | High-Risk Units | [10] |
| **Sleep Disturbances** | 25–36% | General HCW Population | [10] |

These findings suggest a profound and multifaceted psychological burden among healthcare providers during pandemic conditions. Without proper support systems, the persistence of these conditions may affect not only the well-being of HCWs but also the quality and sustainability of healthcare delivery.

**3. Risk and Protective Factors**

Understanding the risk and protective factors that influence the psychological health of healthcare workers (HCWs) during pandemic outbreaks is critical for shaping institutional policies and support mechanisms. This section synthesizes common contributors to psychological distress and highlights mitigating factors that promote resilience and mental well-being.

### 3.1. Risk Factors Contributing to Psychological Distress

Healthcare workers face multiple, often compounding, stressors during pandemics. The following risk factors were consistently reported across studies:

- **Extended Working Hours and Shift Burden:** Long and irregular shifts, often extending beyond 12 hours, were associated with fatigue, reduced coping capacity, and increased rates of depression and anxiety [14]. The lack of adequate rest and work-life balance exacerbated psychological strain, especially among junior staff and residents.
- **Shortages of Personal Protective Equipment (PPE):** Inadequate access to PPE increased fear of infection, especially during the early phases of the COVID-19 pandemic. This fear not only endangered physical safety but also heightened anxiety and reduced morale [15].
- **Social Stigma and Discrimination:** HCWs were often stigmatized by communities and even family members due to perceived contagion risks. Such social rejection contributed to emotional isolation and a sense of alienation [14].
- **Media Overexposure and Misinformation:** Constant exposure to sensationalized news and misinformation created a climate of fear and confusion. Media narratives emphasizing death tolls and system collapse further elevated stress levels [15].
- **Separation from Family and Social Support:** Many frontline workers self-isolated from their families to avoid infecting them. This physical disconnection and emotional detachment significantly impacted their psychological stability [14].

### 3.2. Protective Factors and Coping Mechanisms

Despite the adversities, several protective factors emerged as buffers against mental health deterioration:

- **Organizational Support and Leadership:** Hospitals that implemented transparent communication, flexible work schedules, mental health helplines, and safety protocols observed better psychological outcomes among staff [16]. The perceived fairness and concern from leadership fostered trust and reduced stress.
- **Peer Support and Team Cohesion:** Peer encouragement and a shared sense of purpose helped foster emotional resilience. Team-based debriefings and peer-check-ins were found to be particularly effective in high-stress departments like emergency care [17].
- **Adequate Training and Preparedness:** HCWs who received proper training in infection control and patient management reported lower anxiety levels. Training improved confidence and reduced fear of incompetence or failure under pressure [16].
- **Access to Psychological First Aid (PFA):** On-site mental health professionals and rapid access to counseling services played a significant role in preventing acute stress from progressing to chronic psychological disorders. Structured interventions such as PFA and online cognitive behavioral therapy (CBT) were particularly beneficial [17].

## 4. Regional and Professional Variations

The mental health impacts of pandemics on healthcare workers (HCWs) are not uniformly distributed but vary significantly based on regional context, professional role, and institutional setting. These variations provide critical insights into how socioeconomic conditions, job responsibilities, and organizational environments influence psychological outcomes.

### 4.1. Regional Context: South Asian Vulnerabilities

Healthcare workers in South Asia, particularly in countries like Pakistan, India, and Bangladesh, reported higher levels of psychological distress during pandemics, compared to those in high-income countries. This heightened vulnerability is linked to:

- **Resource Constraints:** Public healthcare facilities in Pakistan often suffer from underfunding, overcrowding, and insufficient access to protective gear, contributing to heightened anxiety and fear among staff [18].
- **Cultural Stigma:** Mental health is often stigmatized in South Asian cultures, deterring HCWs from seeking psychological support. This social barrier leads to underreporting and untreated mental conditions [18].
- **Policy Gaps:** The absence of institutionalized mental health support programs in many South Asian healthcare systems left frontline workers more exposed to burnout and psychological trauma.

### 4.2. Role-Based Differences: Doctors vs. Nurses

Numerous studies revealed profession-specific differences in stress levels and mental health outcomes:

- Nurses reported significantly higher stress and burnout compared to doctors, largely due to prolonged and direct patient care responsibilities, frequent exposure to infection, and limited authority in decision-making [19].
- Nurses also experience greater emotional labor, often acting as mediators between patients and families during critical conditions, particularly when visitation is restricted due to infection control.
- In contrast, doctors reported more decision-related stress tied to medical uncertainty, triage dilemmas, and ethical conflicts in critical care settings [19].

### 4.3 Institutional Settings: Public vs. Private Hospitals

The organizational structure and patient volume in public and private healthcare settings also influenced the psychological well-being of HCWs:

- Public Sector HCWs experienced higher levels of burnout and fatigue, primarily due to higher patient inflow, resource scarcity, and overburdened infrastructure during pandemic peaks [20].
- Private Sector Workers reported lower psychological strain in general, benefitting from better staffing ratios, access to PPE, and organizational support, though some private hospitals faced criticism for delayed crisis response and financial insecurities [20].

**Table: Comparative Mental Health Impact by Role and Institution Type**

| Factor | Doctors | Nurses | Public Hospitals | Private Hospitals |
|---|---|---|---|---|
| **Stress Level** | Moderate to High | High | Very High | Moderate |
| **Burnout Prevalence** | 35–50% | 45–60% | 55–65% | 25–40% |
| **Major Stressors** | Decision fatigue | Patient proximity | Overcrowding, PPE shortage | Organizational instability |
| **Coping Resource Access** | Moderate | Low to Moderate | Low | Moderate to High |

These disparities highlight the need for targeted interventions that account for both professional roles and regional healthcare dynamics. Strengthening mental health infrastructure in resource-constrained settings, reducing workload through equitable task distribution, and encouraging psychological support-seeking—especially among nurses and public sector workers—are crucial strategies to ensure the well-being of the healthcare workforce.

## 5. Policy Recommendations and Interventions

The findings of this systematic review underscore an urgent need to implement structured and proactive mental health policies tailored to support healthcare workers (HCWs) during pandemics. Psychological well-being is foundational not only to individual functioning but also to the resilience and sustainability of health systems during public health emergencies. Based on evidence from the literature, the following recommendations are proposed:

### 5.1. Mandatory Mental Health Screenings

Routine mental health screenings should be integrated into standard occupational health assessments during pandemic periods. These screenings—conducted using validated tools such as the General Health Questionnaire (GHQ) or the Depression Anxiety Stress Scales (DASS)—can facilitate early detection of mental health concerns and trigger timely interventions [21].

### 5.2. Deployment of Mental Health Professionals in Hospitals

Hospitals should ensure the **availability of in-house clinical psychologists and psychiatrists** during health crises. These professionals can conduct individual and group counseling, lead stress management workshops, and provide psychological first aid (PFA) to high-risk HCWs. Embedding mental health experts into infection control and emergency response teams can improve accessibility and reduce stigma [22].

### 5.3. Establishment of 24/7 Psychological Helplines

Dedicated mental health helplines operated by trained psychologists and crisis counselors should be available 24/7 for HCWs. Such services have proven effective in China and Italy during COVID-19 for managing acute stress, anxiety, and suicidal ideation [21]. Confidentiality assurance is critical to encouraging utilization of these services.

### 5.4. Online Cognitive Behavioral Therapy (CBT) and Resilience Training

Given the limitations of in-person therapy during infectious disease outbreaks, online CBT platforms provide a scalable and accessible solution. Studies have shown that internet-based CBT can significantly reduce symptoms of depression and anxiety among HCWs [22]. In parallel, resilience training modules—focusing on adaptive coping, mindfulness, and emotional regulation—can enhance psychological preparedness and buffer stress during crisis periods.

### 5.5. Integration into Disaster Preparedness Protocols

Mental health support should be recognized as a core component of national disaster preparedness and response frameworks. Protocols must include:

- Mental health contingency planning
- Funding for psychological support programs
- Designation of mental health coordinators in emergency task forces
- Collaboration with NGOs and international health organizations [23]

**Box: Key Policy Actions at a Glance**

| Intervention | Description | Stakeholders |
|---|---|---|
| Mandatory Screenings | Early identification of mental health risks | Hospitals, Health Ministries |
| On-Site Mental Health Professionals | Access to immediate support and counseling | Hospital Admin, HR Departments |
| 24/7 Helplines | Crisis support for acute psychological distress | Telehealth Units, NGOs |
| Online CBT and Resilience Training | Digital therapy and preventive coping training | Public Health Authorities, NGOs |
| Disaster Protocol Integration | Embedding mental health into national emergency strategies | NDMA, WHO, Health Commissions |

These recommendations not only aim to address the current gaps observed during COVID-19 and past outbreaks but also serve as a blueprint for mental health resilience in future pandemics. Institutionalizing mental health within healthcare governance can protect frontline workers from long-term psychological harm and foster a more prepared, compassionate, and effective health response system.

**Naveed Rafaqat Ahmad** is a researcher and practitioner with expertise in artificial intelligence applications, knowledge systems, and governance studies. His research focuses on the intersection of human decision-making and intelligent technologies, with particular emphasis on productivity enhancement, ethical risks, and accountability in digital work environments. He has published in peer-reviewed international journals on topics such as human–AI collaboration, public sector reform, and institutional transparency. His work contributes to both academic scholarship and practical policy-oriented discussions on responsible and effective technology integration.

**Figures and Charts**





**Figure 1: Bar Graph – Prevalence of Psychological Disorders Among HCWs During Pandemics**

- X-axis: Mental Health Conditions (Anxiety, Depression, PTSD, Insomnia, Burnout)
- Y-axis: Prevalence Percentage (%)
- COVID-19 shows highest values (e.g., Anxiety: 45%, Depression: 30%)





**Figure 2: Line Graph – Mental Health Incidence Among HCWs Over Time (2003–2023)**

- X-axis: Year
- Y-axis: Number of Reported Psychological Cases per 1,000 HCWs
- Spikes during SARS (2003), MERS (2012), Ebola (2014), and COVID-19 (2020–2022)

## Summary

The systematic review highlights the critical psychological toll pandemics have on healthcare workers globally, with significant implications for the Pakistani context. Mental health challenges such as anxiety, depression, PTSD, and burnout were consistent across pandemics, though exacerbated during COVID-19 due to prolonged duration and intensity. Risk factors like inadequate PPE, social isolation, and overwork, particularly in South Asia, call for robust interventions. Policymakers must institutionalize mental health support mechanisms as a part of health emergency responses. Interdisciplinary strategies including counseling, organizational reform, and tele-mental health platforms are necessary to safeguard HCW well-being in future outbreaks.

## References

WHO. (2021). Mental health and psychosocial considerations during COVID-19 outbreak.

Greenberg N, et al. (2020). Managing mental health challenges faced by HCWs during COVID-19. BMJ.

Brooks SK, et al. (2020). The psychological impact of quarantine. Lancet.

Maunder R, et al. (2006). Long-term psychological effects of SARS. Emerg Infect Dis.

Lai J, et al. (2020). Factors associated with mental health outcomes among HCWs exposed to COVID-19. JAMA Netw Open.

Pappa S, et al. (2020). Prevalence of depression, anxiety, and insomnia among HCWs. Brain Behav Immun.

Moher D, et al. (2009). Preferred reporting items for systematic reviews and meta-analyses: PRISMA statement. PLoS Med.

Critical Appraisal Skills Programme (CASP). (2023). Checklist for systematic reviews.

Shaukat N, et al. (2020). Physical and mental health impacts of COVID-19 on healthcare workers: A scoping review. Int J Emerg Med.

Rajkumar RP. (2020). COVID-19 and mental health: A review. Asian J Psychiatr.

Spoorthy MS, et al. (2020). Mental health problems faced by HCWs due to COVID-19 pandemic. Asian J Psychiatr.

Barello S, et al. (2020). Burnout among HCWs during COVID-19. J Health Soc Sci.

Rossi R, et al. (2020). Mental health outcomes among HCWs in Italy. JAMA Netw Open.

Chew NWS, et al. (2020). Psychological impact of the COVID-19 pandemic on healthcare workers. Ann Intern Med.

Khan A, et al. (2021). Mental health of Pakistani frontline healthcare workers. Pak J Med Sci.
Zhang SX, et al. (2020). Protecting mental health of HCWs. J Occup Environ Med.
Walton M, et al. (2020). Mental health care for medical staff in China during the COVID-19 outbreak. Lancet Psychiatry.
Khalid I, et al. (2020). Psychological distress and resilience among medical staff in Pakistan. Pak J Public Health.
Shechter A, et al. (2020). Psychological distress in healthcare workers in NYC during the pandemic. Gen Hosp Psychiatry.
Abbasi J. (2020). The impact of pandemic on healthcare delivery. JAMA.
Ho CS, et al. (2020). Interventions to address mental health of HCWs during pandemics. Psychiatry Res.
Liu S, et al. (2020). Online mental health services in China during COVID-19 outbreak. Lancet Psychiatry.
IASC. (2020). Guidelines on mental health and psychosocial support in emergency settings.
Ahmad, N. R. (2024). *Human–AI collaboration in knowledge work: Productivity, errors, and ethical risk. Journal of Knowledge Systems and Digital Ethics, 6*(2), Article 9250. https://doi.org/10.52152/6q2p9250